

## Math 205 Handout 3: Induction and Pigeonholes

- **Induction:** Given a statement  $p(n)$  about an integer  $n$  we wish to show it is true for all integer values of  $n$  at least  $a$  and we proceed as follows:

- *Initial Case:* Show that  $p(a)$  is true (optionally also test  $p(a+1)$  and  $p(a+2)$  to see how the induction will proceed).
- *Inductive Case:* Assume  $p(k)$  is true for some value of  $k \geq a$ . State one side of  $p(k+1)$  in terms of the corresponding side of  $p(k)$  and use the assumptions to deduce that the other side of  $p(k+1)$  is related in the same way as  $p(n)$  was.

For example:  $p(n) := \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$ ,

- *Initial Case:* The first possible value of  $n$  is 1, so we consider  $p(1) := "1^2 = 1 = \frac{1 \times (1+1) \times (2 \times 1 + 1)}{6} = 1"$  as required. Similarly,  $p(2) := "1^2 + 2^2 = 5 = \frac{2 \times (2+1) \times (2 \times 2 + 1)}{6} = 5"$  and  $p(3) := "1^2 + 2^2 + 3^2 = 14 = \frac{3 \times (3+1) \times (2 \times 3 + 1)}{6} = 14"$ .
- *Inductive Case:* Assume  $p(k) := \sum_{i=1}^k i^2 = \frac{k(k+1)(2k+1)}{6}$ . Now the left hand side of  $p(k+1)$  is

$$\sum_{i=1}^{k+1} i^2 = \left( \sum_{i=1}^k i^2 \right) + (k+1)^2 = \text{LHS}(p(k)) + (k+1)^2.$$

But using the assumption (the inductive hypothesis), we get that

$$\begin{aligned} \sum_{i=1}^{k+1} i^2 &= \frac{k(k+1)(2k+1)}{6} + (k+1)^2 \\ &= (k+1) \frac{k(2k+1) + 6(k+1)}{6} \\ &= (k+1) \frac{(2k^2 + 7k + 6)}{6} \\ &= (k+1) \frac{(2k+3)(k+2)}{6} \\ &= \frac{(k+1)(k+2)(2k+3)}{6} \\ &= \frac{(k+1)((k+1)+1)(2(k+1)+1)}{6} \end{aligned}$$

But this is exactly the statement  $p(k+1)$  that we wished to establish!

- **Basic formulae in Sigma Notation:**

$$1 + \dots + n = \sum_{i=1}^n i = \frac{n(n+1)}{2} \quad b + \dots + b = \sum_{i=1}^n b = nb \quad 1 + x + \dots + x^n = \sum_{i=0}^n x^i = \frac{x^{n+1} - 1}{x - 1}$$

- **Pigeonhole Principle:**

Given a set of  $n$  objects and  $m$  groups to place them in, if  $m < n$  then at least one group has two objects in it. In general, we can say that one group must have at least

$$\left\lfloor \frac{n-1}{m} \right\rfloor + 1$$

objects in it where the brackets round down the nearest integer.