

## Math205 Handout 2: Methods of Proof

We shall show how to go about proving the statement “if an odd integer is multiplied by -1 and that new integer is then added to 26 the result is odd”. We first identify the statements  $p(x)$  and  $q(x)$  in the above statement if it is  $p(x) \rightarrow q(x)$  and deduce that:

$$p(x) \quad \equiv \quad “x \text{ is odd}” \quad \equiv \quad “x = 2j + 1 \text{ for some } j \in \mathbb{Z}”$$

$$q(x) \quad \equiv \quad “26 - x \text{ is odd}” \quad \equiv \quad “26 - x = 2k + 1 \text{ for some } k \in \mathbb{Z}”$$

We are usually either told to use one of the methods below, or we can choose one:

- **Direct:** We suppose that  $p(x)$  is true and using what that tells us about  $x$  we then apply that to the subject of  $q(x)$  in order to try to show that it is true when  $p(x)$  is.

So if  $p(x)$  is true then  $x = 2j + 1$ , and  $q(x)$  is about  $26 - x$ , and putting these two things together

$$26 - x = 26 - (2j + 1) = 26 - 2j - 1 = 25 - 2j = 25 + 2 \times (-j) = 1 + 24 + 2 \times (-j) = 2 \times (12 - j) + 1$$

Thus we have our statement in the form of  $q(x)$  where  $k = 12 - j$ , and it just remains to establish that this  $k$  is an integer. Since  $j$  is, multiplying by -1 means  $-j$  is still an integer, and then subtracting this from 12, another integer, means that  $k$  is an integer as required.

- **Contrapositive:** We can alternatively suppose that  $q(x)$  is false and using what that tells us about  $x$  we then apply that to the subject of  $p(x)$  in order to try to show that it is false when  $p(x)$  is. We are thus proving  $(\sim q(x)) \rightarrow (\sim p(x))$  which we know is logically equivalent to  $p(x) \rightarrow q(x)$ .

So if  $q(x)$  is false then  $26 - x = 2m, m \in \mathbb{Z}$ , and  $\sim p(x)$  says that “ $x = 2n, n \in \mathbb{Z}$ ”. Simplifying  $(\sim q(x))$  to tell us about  $x$

$$\begin{aligned} 26 - x &= 2m \\ x &= 26 - 2m \\ &= 2 \times (13 - m) \end{aligned}$$

This equals  $2n$  if we take  $n = 13 - m$  and so, again, since  $m$  is an integer, so is  $-m$  and adding 13 to this keeps it an integer, so  $n$  is an integer as required.

- **Contradiction:** We now suppose that  $p(x)$  is true and also that  $q(x)$  is false. We intend to get an impossible situation arising whence we can use the logical equivalence of  $(p(x) \wedge (\sim q(x))) \leftrightarrow (\sim T_0)$  and  $(p(x) \rightarrow q(x)) \leftrightarrow T_0$  to show that  $p(x)$  implies  $q(x)$  as required.

As before, if  $p(x)$  is true then  $x = 2j + 1$ , and  $(\sim q(x))$  says that  $26 - x = 2m$ . Combining these two statements to remove  $x$  we get:

$$\begin{aligned} 26 - (2j + 1) &= 2m \\ 25 &= 2j + 2m \\ &= 2(j + m) \\ \frac{25}{2} &= j + m \end{aligned}$$

This statement is our desired contradiction since both  $j$  and  $m$  are integers and so their sum is an integer, but  $\frac{25}{2}$  is certainly not an integer as it is 12.5 in decimal terms and no integer has to be written with a decimal point.